

Abstract. Simple formulas are given for generating Chern-Simons basic invariant polynomials by repeated exterior differentiation for n -dimensional differentiable manifolds having a general linear connection.

PACS numbers: 02.40.-k, 04.20.Fy

This paper presents simple formulas for generating Chern-Simons basic invariant polynomials by repeated exterior differentiation for n -dimensional differentiable manifolds having a general linear connection.

The Chern-Simons basic invariant polynomials of such a manifold M are the $(2p-1)$ -forms $Q_p(\omega \wedge \Omega^{p-1})$ together with the $2p$ -forms $Q_p(\Omega^p)$ defined, for $p > 0$, by¹

$$\begin{aligned} Q_p(\omega \wedge \Omega^{p-1}) &= \omega_{i_p}^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{p-3}}^{i_{p-2}} \wedge \Omega_{i_{p-2}}^{i_{p-1}} \wedge \Omega_{i_{p-1}}^{i_p} \\ &= \text{tr}(\omega \wedge \Omega^{p-1}) \end{aligned} \quad (1)$$

and

$$\begin{aligned} Q_p(\Omega^p) &= \Omega_{i_p}^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{p-3}}^{i_{p-2}} \wedge \Omega_{i_{p-2}}^{i_{p-1}} \wedge \Omega_{i_{p-1}}^{i_p} \\ &= \text{tr}(\Omega^p), \end{aligned} \quad (2)$$

respectively, where ω_a^b and Ω_a^b are the connection 1-forms and curvature 2-forms, respectively, of M .

The 1st ordinary exterior differentials of the basis tangent vectors $\mathbf{e}_a$ of M are given by²⁻⁴

$$d\mathbf{e}_a = \mathbf{e}_b \omega_a^b, \quad (3)$$

the contractions of which with the basis 1-forms ω^b of M are given by

$$\langle \omega^b, d\mathbf{e}_a \rangle = \omega_a^b \quad (4)$$

and in view of which the 1st absolute exterior differentials of $\mathbf{e}_a$ are given by

$$\begin{aligned} D\mathbf{e}_a &= d\mathbf{e}_a - \mathbf{e}_b \omega_a^b \\ &= 0, \end{aligned} \quad (5)$$

where the contractions of ω^b with $\mathbf{e}_a$ are given by

$$\langle \omega^b, \mathbf{e}_a \rangle = \delta_a^b, \quad (6)$$

where δ_a^b is the Kronecker delta.

The 2nd ordinary exterior differentials of $\mathbf{e}_a$ are given by⁵⁻¹²

$$\begin{aligned} d^2\mathbf{e}_a &= d d\mathbf{e}_a \\ &= d\mathbf{e}_b \omega_a^b \\ &= (d\mathbf{e}_b) \wedge \omega_a^b + \mathbf{e}_b d\omega_a^b \\ &= \mathbf{e}_c \omega_b^c \wedge \omega_a^b + \mathbf{e}_b d\omega_a^b \\ &= \mathbf{e}_b (d\omega_a^b + \omega_c^b \wedge \omega_a^c) \\ &= \mathbf{e}_b \Omega_a^b, \end{aligned} \quad (7)$$

the contractions of which with ω^b are given by

$$\langle \omega^b, d^2\mathbf{e}_a \rangle = \Omega_a^b \quad (8)$$

and where

$$\begin{aligned} \Omega_a^b &= d\omega_a^b + \omega_c^b \wedge \omega_a^c \\ &= D\omega_a^b + \omega_a^c \wedge \omega_c^b \\ &= d\omega_a^b - \omega_a^c \wedge \omega_c^b \\ &= D\omega_a^b - \omega_c^b \wedge \omega_a^c. \end{aligned} \quad (9)$$

The 3rd ordinary exterior differentials of $\mathbf{e}_a$ are given by

$$\begin{aligned} d^3\mathbf{e}_a &= d d^2\mathbf{e}_a \\ &= d\mathbf{e}_b \Omega_a^b \\ &= (d\mathbf{e}_b) \wedge \Omega_a^b + \mathbf{e}_b d\Omega_a^b \\ &= (\mathbf{e}_c \omega_b^c) \wedge \Omega_a^b + \mathbf{e}_b d\Omega_a^b \\ &= \mathbf{e}_c \Omega_b^c \wedge \omega_a^b \end{aligned} \quad (10)$$

using Bianchi's identity for Ω_a^b ,¹³ i.e.,

$$\begin{aligned} D\Omega_a^b &= d\Omega_a^b - \omega_a^c \wedge \Omega_c^b + \omega_c^b \wedge \Omega_a^c \\ &= 0, \end{aligned} \quad (11)$$

as well as by

$$\begin{aligned} d^3\mathbf{e}_a &= d^2 d\mathbf{e}_a \\ &= d^2\mathbf{e}_b \omega_a^b \\ &= (d^2\mathbf{e}_b) \wedge \omega_a^b + \mathbf{e}_b d^2\omega_a^b \\ &= (\mathbf{e}_c \Omega_b^c) \wedge \omega_a^b + 0 \\ &= \mathbf{e}_c \Omega_b^c \wedge \omega_a^b \end{aligned} \quad (12)$$

¹ Chern, S. S., and J. Simons, "Characteristic forms and geometric invariants," *Annals of Math.*, **99** (1974) 48.

² Misner, C. W., K. S. Thorne, and J. A. Wheeler, *Gravitation*, W. H. Freeman & Co., San Francisco, CA (1973), p. 350.

³ Cartan, É. J., *Leçons sur la géométrie des espaces de Riemann*, Gauthier-Villars et C^{ie}, Paris, France (1928), p. 180; *Leçons sur la géométrie des espaces de Riemann*, 2nd édition, rev. et augm., Gauthier-Villars, Paris, France (1946), p. 180.

⁴ Chern, S. S., "A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds," *Ann. of Math.*, **45** (1944) 747; "On the curvatura integra in a Riemannian manifold," *ibid.*, **46** (1945) 674; "On the Minimal Immersions of the Two-sphere in a Space of Constant Curvature," in Gunning, R. C. (ed.), *Problems in Analysis: A Symposium in Honor of Salomon Bochner*, Princeton University Press, Princeton, NJ (1970), p. 27.

⁵ Bott, R., and S. S. Chern, "Some Formulas Related to Complex Transgression," in Haefliger, A., and R. Narasimhan (eds.), *Essays on Topology and Related Topics; Memoires dédiés à Georges de Rham*, Springer-Verlag, New York, NY (1970), p. 48.

⁶ Cartan, É. J., "Sur les variétés à connexion affine et la théorie de la relativité généralisée (Première Partie)," *Annales scientifiques de l'École Normale supérieure*, **40** (1923) 325; (1928), *op. cit.*, p. 182; (1946), *op. cit.*, p. 181.

⁷ Chern, S. S., "Differential Geometry of Fiber Bundles," in *Proceedings of the International Congress of Mathematicians*, **2** (1950) 397; "Topics in Differential Geometry," [mimeographed notes], The Institute for Advanced Study, Princeton, NJ (1951), Chap. 3, Sec. 3; (1970), *op. cit.*

⁸ Cohen, J. M., *Exterior Differential Forms and Orthonormal Cartan Frames*, [unpublished monograph], (1975), p. 33.

⁹ Flanders, H., "Development of an Extended Exterior Differential Calculus," *Trans. Amer. Math. Soc.*, **75** (1953) 311; "Methods in Affine Connection Theory," *Pac. J. Math.*, **5** (1955) 391.

¹⁰ Misner, C. W., et al., *op. cit.*, pp. 350-351 and 359.

¹¹ Petrie, T., and J. Randall, *Connections, Definite Forms, and Four-manifolds*, Clarendon Press, Oxford, England (1990), pp. 11-12.

¹² Trautmann, A., *Differential Geometry for Physicists—Stony Brook Lectures*, Bibliopolis, edizioni di filosofia e scienze, Napoli, Italy (1984), pp. 91-92.

¹³ Cartan, É. J., (1928), *op. cit.*, p. 214; (1946), *op. cit.*, p. 210.

using Poincaré's theorem for scalar-valued exterior differential forms,¹⁴⁻¹⁵

i.e.,

$$d^2 \alpha = 0, \quad (13)$$

where α is an arbitrary scalar-valued exterior differential form.

The 4th ordinary exterior differentials of e_a are given by

$$\begin{aligned} d^4 e_a &= d^2 d^2 e_a \\ &= d^2 e_b \Omega_a^b \\ &= (d^2 e_b) \wedge \Omega_a^b + e_b d^2 \Omega_a^b \\ &= (e_c \Omega_b^c) \wedge \Omega_a^b + 0 \\ &= e_c \Omega_b^c \wedge \Omega_a^b. \end{aligned} \quad (14)$$

In general, the p^{th} ordinary exterior differentials of e_a for $p > 0$ are given by

$$\begin{aligned} d^p e_a &= \begin{cases} e_{i_1} \Omega_{i_2}^{i_1} \wedge \Omega_{i_3}^{i_2} \wedge \Omega_{i_4}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-1)/2}}^{i_{(p-2)/2}} \wedge \Omega_{i_{(p+1)/2}}^{i_{(p-3)/2}} \wedge \omega_a^{i_{(p+1)/2}}, & \text{if } p \text{ is odd} \\ e_{i_1} \Omega_{i_2}^{i_1} \wedge \Omega_{i_3}^{i_2} \wedge \Omega_{i_4}^{i_3} \wedge \dots \wedge \Omega_{i_{p/2}}^{i_{(p-4)/2}} \wedge \Omega_{i_{p/2}}^{i_{(p-2)/2}} \wedge \Omega_a^{i_{p/2}}, & \text{if } p \text{ is even} \end{cases} \\ &= \begin{cases} e_{i_1} \omega_a^{i_{(p+1)/2}} \wedge \Omega_{i_{(p+1)/2}}^{i_{(p-1)/2}} \wedge \Omega_{i_{(p-1)/2}}^{i_{(p-3)/2}} \wedge \dots \wedge \Omega_{i_4}^{i_3} \wedge \Omega_{i_3}^{i_2} \wedge \Omega_{i_2}^{i_1}, & \text{if } p \text{ is odd} \\ e_{i_1} \Omega_a^{i_{p/2}} \wedge \Omega_{i_{p/2}}^{i_{(p-2)/2}} \wedge \Omega_{i_{(p-2)/2}}^{i_{(p-4)/2}} \wedge \dots \wedge \Omega_{i_4}^{i_3} \wedge \Omega_{i_3}^{i_2} \wedge \Omega_{i_2}^{i_1}, & \text{if } p \text{ is even} \end{cases} \\ &= \begin{cases} e_{i_{(p+1)/2}} \omega_a^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-5)/2}}^{i_{(p-3)/2}} \wedge \Omega_{i_{(p-3)/2}}^{i_{(p-1)/2}} \wedge \Omega_{i_{(p-1)/2}}^{i_{(p+1)/2}}, & \text{if } p \text{ is odd} \\ e_{i_{p/2}} \Omega_a^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-6)/2}}^{i_{(p-4)/2}} \wedge \Omega_{i_{(p-4)/2}}^{i_{(p-2)/2}} \wedge \Omega_{i_{(p-2)/2}}^{i_{p/2}}, & \text{if } p \text{ is even} \end{cases}, \end{aligned} \quad (15)$$

the contractions of which with ω^a are given by

$$\begin{aligned} \langle \omega^a, d^p e_a \rangle &= \begin{cases} \delta_{i_{(p+1)/2}}^a \omega_a^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-5)/2}}^{i_{(p-3)/2}} \wedge \Omega_{i_{(p-3)/2}}^{i_{(p-1)/2}} \wedge \Omega_{i_{(p-1)/2}}^{i_{(p+1)/2}}, & \text{if } p \text{ is odd} \\ \delta_{i_{p/2}}^a \Omega_a^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-6)/2}}^{i_{(p-4)/2}} \wedge \Omega_{i_{(p-4)/2}}^{i_{(p-2)/2}} \wedge \Omega_{i_{(p-2)/2}}^{i_{p/2}}, & \text{if } p \text{ is even} \end{cases} \\ &= \begin{cases} \omega_{i_{(p+1)/2}}^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-5)/2}}^{i_{(p-3)/2}} \wedge \Omega_{i_{(p-3)/2}}^{i_{(p-1)/2}} \wedge \Omega_{i_{(p-1)/2}}^{i_{(p+1)/2}}, & \text{if } p \text{ is odd} \\ \Omega_{i_{p/2}}^{i_1} \wedge \Omega_{i_1}^{i_2} \wedge \Omega_{i_2}^{i_3} \wedge \dots \wedge \Omega_{i_{(p-6)/2}}^{i_{(p-4)/2}} \wedge \Omega_{i_{(p-4)/2}}^{i_{(p-2)/2}} \wedge \Omega_{i_{(p-2)/2}}^{i_{p/2}}, & \text{if } p \text{ is even} \end{cases} \\ &= \begin{cases} Q_{(p+1)/2}(\omega \wedge \Omega^{(p-1)/2}), & \text{if } p \text{ is odd} \\ Q_{p/2}(\Omega^{p/2}), & \text{if } p \text{ is even} \end{cases}. \end{aligned} \quad (16)$$

Equation (16) then yields the formulas in question, viz.,

$$Q_p(\omega \wedge \Omega^{p-1}) = \langle \omega^a, d^{2p-1} e_a \rangle \quad (17)$$

and

$$Q_p(\Omega^p) = \langle \omega^a, d^{2p} e_a \rangle. \quad (18)$$

For an aside, note that using Eq. (18) and taking the 0th ordinary exterior differentials of e_a to be given by

$$d^0 e_a = e_a, \quad (19)$$

the polynomial $Q_p(\Omega^p)$ for $p = 0$ is given by

$$\begin{aligned} Q_0(\Omega^0) &= \langle \omega^a, d^0 e_a \rangle \\ &= \langle \omega^a, e_a \rangle \\ &= \delta_a^a \\ &= n, \end{aligned} \quad (20)$$

where n , having appeared above, is the number of dimensions of M .

¹⁴ Cartan, É. J., (1928), *op. cit.*, pp. 209 and 215; *Le systèmes différentiels extérieurs et leurs applications géométriques*, Actualités scientifiques et industrielles, 994, Exposés de géométrie, vol. 12, Librairie scientifiques Hermann & C^{ie}, Éditeurs, Paris, France (1945), p. 37; (1946), *op. cit.*, pp. 208 and 211.

¹⁵ Chern, S. S., "The Geometry of Isotropic Surfaces," *Annals of Math.*, **43** (1942) 545; "On the Euclidean Connections in a Finsler Space," *Proc. Nat. Acad. Sci.*, **29** (1943) 33; "Local Equivalence and Euclidean Connections in Finsler Spaces," *Science Reports Nat. Tsing Hua Univ.*, **5** (1948) 95.